

List of REAP Publications (December 2020)

- [92] Lin, S.-K., Yang, S.-Y., Park, S.-C., Jang, O.-J., Zhu, X., Xiang, Y.-T., Ouyang, W.-C., Javed, A., Khan, M.N.S., Grover, S., Avasthi, A., Kallivayalil, R.A., Chee, K.Y., Chami, N., Kato, T.A., Hayakawa, K., Pariwatcharakul, P., Maramis, M., Seneviratne, L., Kang, S., Tang, W.K., Oo Oo, T., Sartorius, N., Tan, C.-H., Chong, M.-Y., Park, Y.C., Shinfuku, N. (2020). Prescription Patterns for Bipolar Disorder in Asian Countries: Findings from REAP-BD. *Clinical Psychopharmacology and Neurosciences*. Published online December 26, 2020.
- [91] Xu, S.-W., Dong, M., Zhang, Q., Yang, S.-Y., Chen, L.-Y., Sim, K., He, Y.-L., Chiu, F.K.H., Sartorius, N., Tan, C.-H., Chong, M.-Y., Shinfuku, N., Lin, S.-K., Ng, C.H., Ungvari, G.S., Najooan, E., Kallivayalil, R.A., Jamaluddin, R., Javed, A., Iida, H., Swe, T., Zhang, B., Xiang, Y.-T. (2020, May). Clozapine prescription pattern in patients with schizophrenia in Asia: The REAP survey (2016). *Psychiatry Research*, 287:112271–112271
- [90] Park, S.-C., Jang, E.Y., Xiang, Y.-T., Kanba, S., Kato, T.A., Chong, M.Y., Lin, S.K., Yang, S.-Y., Avasthi, A., Grover, S., Kallivayalil, R.A., Udomratn, P., Chee, K.Y., Tanra, A.J., Tan, C.-H., Sim, K., Sartorius, N., Park, Y.C., Shinfuku, N. (2020, February 12). Network analysis of the depressive symptom profiles in Asian patients with depressive disorders: Findings from the Research on Asian Psychotropic Prescription Patterns for Antidepressants (REAP-AD). *Psychiatry and Clinical Neurosciences*.
- [89] Lin, S.-K. Antipsychotic Polypharmacy: A Dirty Little Secret or a Fashion? (2020, February). *Int J Neuropsychopharmacol*, 23(2),125–131.
- [88] Malik, M., Ghani, M.U., Mazhar, W., Munir, N. (2019, May-June). Changing trends in anti-psychotic prescription pattern in Pakistan. *Pak J Med Sci*, 35(3).
- [87] Dong, M., Zeng, L.N., Zhang, Q., Yang, S.Y., Chen, L.Y., Najooan, E., Kallivayalil, R.A., Viboonma, K., Jamaluddin, R., Javed, A., Hoa, D.T.Q., Iida, H., Sim, K., Swe, T., He, Y.L., Park, Y., Ahmed, H.U., De Alwis, A., Chiu, H.F.K., Sartorius, N., Tan, C.H., Chong, M.Y., Shinfuku, N., Lin, S.K., Avasthi, A., Grover, S., Ng, C.H., Ungvari, G.S., Xiang, Y.T. (2019, October). Prescription of antipsychotic and concomitant medications for adult Asian schizophrenia patients: Findings of the 2016 Research on Asian Psychotropic Prescription Patterns (REAP) survey. *Asian J Psychiatry*, 45, 74-80, <https://doi.org/10.1016/j.ajp.2019.08.010>
- [86] Dong, M., Zeng, N., Zhang, Q., Yang, S.Y., Chen, L.Y., Sim, K., He, Y.L., Chiu, H.F., Sartorius, N., Tan, C.H., Chong, M.Y., Shinfuku, N., Lin, S.K., Ng, C.H., Ungvari, G.S., Xiang, Y.T. (2019, November). Antipsychotic Polypharmacy in Older Adult Asian Patients With Schizophrenia: Research on Asian Psychotropic Prescription Pattern. *J Geriatr Psychiatry*

Neurol, 32(6), 304-311.

<https://doi.org/10.1177/0891988719862636>

[85] Dong, M., Zeng, L.N., Zhang, Q., Ungvari, G.S., Ng, C.H., Chiu, H.F.K., Si, T.M., Sim, K., Avasthi, A., Grover, S., Chong, M.Y., Chee, K.Y., Kanba, S., Lee, M.S., Yang, S.Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Mahendran, R., Tan, C.H., Shinfuku, N., Xiang, Y.T. (2019, July). Concurrent antipsychotic use in older adults treated with antidepressants in Asia. *Psychogeriatrics*, 19(4), 333-339.

<https://doi.org/10.1111/psyg.12416>

[84] Zhong, X.M., Wang, F., Zhang, Q., Ungvari, G.S., Ng, C.H., Chiu, H.F.K., Si, T.M., Sim, K., Avasthi, A., Grover, S., Chong, M.Y., Chee, K.Y., Kanba, S., Lee, M.S., Yang, S.Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Mahendran, R., Tan, C.H., Shinfuku, N., Xiang, Y.T. (2019, May). Concurrent benzodiazepine use in older adults treated with antidepressants in Asia. *Int Psychogeriatr*, 31(5), 685-691

<https://doi.org/10.1017/S1041610217002563>

[83] Park, S.-C., Jang, E.Y., Xiang, Y.-T., Kanba, S., Kato, T.A., Chong, M.-Y., Lin, S.-K., Yang, S.-Y., Avasthi, A., Grover, S., Kallivayalil, R.A., Udomratn, P., Chee, K.Y., Tanra, A.J., Tan, C.-H., Sim, K., Sartorius, N., Park, S.C., Shinfuku, N. Network Analysis of the Depressive Symptom Profiles in Asian Patients with Depressive Disorders: Findings from the Research on Asian Psychotropic Prescription Patterns for Antidepressant (REAP-AD). (2020). *Psychiatry Clin Neurosci* (accepted for publication).

<http://onlinelibrary.wiley.com/doi/10.1111/pcn.12989/full>

[82] Park, S.-C., Jang, E.Y., Kim, K., Lee, H., Choi, J., Dan, A., Hussain, A., Tanra, A.J., Kato, T.A., Chee, K.Y., Lin, S.-K., Javed, A., Tan, C.H., Sartorius, N., Shinfuku, N., Park, Y.C.: Establishing the Cut-off Scores for the Severity Ranges of Schizophrenia on the BPRS-6 Scale: Findings from the REAP-AP. (2019). *Psychiatry Clin Psychopharmacology*, 29(4), 895-898.

<https://doi.org/10.1080/24750573.2019.1695994>

[81] Park, S.-C., Oh, H.S., Tripathi, A., Kallivayalil, R.A., Avasthi, A., Grover, S., Tanra, A.J., Kanba, S., Kato, T.A., Inada, T., Chee, K.Y., Chong, M.-Y., Lin, S.-K., Sim, K., Xiang, Y.T., Tan, C.H., Javed, A., Sartorius, N., Shinfuku, A., Park, Y.C.: Cannabis Use Correlates with Aggressive Behavior and Long-acting Injectable Antipsychotic Treatment in Asian Patients with Schizophrenia. (2019). *Nord J Psychiatry*, 73(6), 323-330.

<https://doi.org/10.1080/08039488.2019.1632381>

[80] Park, S.-C., Avasthi, A., Grover, S., Tanra, A.J., Kato, T.A., Chee, K.Y., Chong, M.-Y., Jang, E.Y., Kim, G.-M., Javed, A., Tan, C.H., Sartorius, N., Shinfuku, N., Park, Y.C. (2019). Establishing the Cut-off Score for Aggression on the Brief Psychiatric Rating Scale-Excited Component (BPRS-EC) in Schizophrenia Patients. *Psychiatr Clin Psychopharmacol*, 29(4), 407-410.

<https://doi.org/10.1080/24750573.2019.1626067>

[79] Park, S.-C., Kim, K., Jang, O.-J., Kim, S.-G., Lee, J.G., Park, J.H., Choi, J., Lee, D.-W., Lin, S.-K., Tan, C.H., Shinfuku, N., Park, Y.C. (2019). Clinical Characteristics and Psychotropic Drug Prescription Patterns of Bipolar Disorder Patients with a History of Suicidal Attempts: Findings from the REAP-BD, Korea. *Psychiatry Investig*, 16(6), 459-463.

<https://doi.org/10.30773/pi.2019.03.10>

[Full text](#)

[78] Kim, K., Yang, H., Na, E., Lee, H., Jang, O.-J., Yoon, H.-J., Oh, H.S., Ham, B.-J., Park, S.-C., Lin, S.-K., Tan, C.H., Shinfuku, N., Park, Y.C. (2019). Examining Patterns of Polypharmacy in Bipolar Disorder: Findings from the REAP-BD, Korea. *Psychiatry Investig*, 16(5), 397-402.

<https://doi.org/10.30773/pi.2019.02.26.4>

[Full text](#)

[77] Xu S-W, Dong M, Zhang Q, Yang S-Y, Chen L-Y, Sim K, He Y-L, Chiu H F-K, Sartorius N, Tan C-H, Chong M-Y, Shinfuku N, Lin S-K, Ng CH, Ungvari GS, Najooan E, Kallivayalil RA, Jamaluddin R, Javed A, Iida H, Swe T, Zhang B, Xiang Y-T. Clozapine prescription pattern in patients with schizophrenia in Asia: The REAP survey (2016). (2019, March 15). *Psychiatry Research*, S0165-1781(18)32218-2. doi: 10.1016/j.psychres.2019.02.056.

[Full text](#)

[76] Dong M, Zeng L-N, Zhang Q, Yang S-Y, Chen L-Y, Sim K, He Y-L, Chiu H F-K, Sartorius N, Tan C-H, Mian-Chong Y, Shinfuku N, Lin S-K, Ng CH, Ungvari GS, Xiang Y-T. Antipsychotic polypharmacy in Asian older adult patients with schizophrenia: Research on Asian Psychotropic Prescription Pattern for Antipsychotics (2016 Data). *Journal of Geriatric Psychiatry and Neurology*.

[75] Dong, M., Zeng, L.-N., Zhang, Q., Ungvari, G.S., Ng, C.H., Chiu, H.F.K, Si, T.-M., Sim, K., Avasthi, A., Grover, S., Chong, M.-Y., Chee, K.-Y., Kanba, S., Lee, M.-S., Yang, S.-Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Mahendran, R., Tan, C.-H., Shinfuku, N., Xiang Y.-T. (2019, February). Concurrent antipsychotics use in older adults treated with antidepressants in Asia. *Psychogeriatrics*, doi: 10.1111/psyg.12416.

[Full text](#)

[74] Park, Y.C., Lee, M.-S., Si, T.-M., Chiu, H.F.K., Kanba, S., Chong, M.-Y., Tripathi, A., Udomratn, P., Chee, K.Y., Tanra, A.J., Rabbani, G., Javed, A., Kathiarachchi, S., Aun, W., Myint, Cuong, T.V., Sim, K., Yang, S.-Y., Sartorius, N., Park, S.-C. (2018, November). Psychotropic Drug-Prescribing Correlates of Disorganized Speech in Asians with Schizophrenia: The REAP-AP Study. *Saudi Pharmaceutical Journal*.

<https://doi.org/10.1016/j.jsps.2018.11.006>

[Full text](#)

[73] Park, Y.-C., Yang, S.-Y., Chong, M.-Y., Kanba, S., Sartorius, N., Shinfuku, N., Tan, C.-H., Park, S.-C. (2018, October). Differences in High Dose Antipsychotic Prescriptions in Patients with Schizophrenia in Asian Countries/Areas: Findings from the REAP-AP Study. *Psychiatry Investig.*, 15(10), 1007–1008.

Published online 2018 Oct 22. doi: [10.30773/pi.2018.09.06]

PMCID: PMC6212696 , PMID: 30373360

[full text](#)

[72] Huang, C.-Y., Yang, S.-Y., Mojtabai, R., Lin, S.-K., He, Y.-L., Chong, M.-Y., Ungvari, G., Tan, C.-H., Xiang, Y.-T., Sartorius, N., Shinfuku, N., Chen, L.-Y. (2018, December). Trends of Polypharmacy and Prescription Patterns of Antidepressants in Asia. *Journal of Clinical Psychopharmacology*, 38(6), 598–603.

[Full text](#)

[71] Zhong, X.-M., Dong, M., Wang, F., Zhang, Q., Ungvari, G.S., Ng, C.H., Chiu, H.F.K, Si, T.-M., Sim, K., Avasthi, A., Grover, S., Chong, M.-Y., Chee, K.-Y., Kanba, S., Lee, M.-S., Yang, S.-Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Mahendran, R., Tan, C.H., Shinfuku, T.N., Xiang, Y.-T. (2018). Physical comorbidities in older adults receiving antidepressants in Asia: Jul 2018, *Psychogeriatrics*. First published: 10 July 2018 <https://doi.org/10.1111/psyg.12327>

[full text](#)

[70] Malik, M., Ghani, U., Awais, N., Munir, N., Javed, A., Hamirani, M., Kahn, N.S., Yang, S.-Y., Chen, L.-Y., Lin, S.K., Sartorius, N., Tan, C.-H., Chong, M.-Y., Shinfuku, M., Grover, S., Shen, W.W., Shinfuku, N. (2018). More than Half of Patients with Schizophrenia Are Receiving Polypharmacy and Co-Prescription of Anxiolytics in Pakistan— Findings from Research on Asian Prescription Pattern in 2016. *Open Journal of Psychiatry*, 8, 199-211.

<https://doi.org/10.4236/ojpsych.2018.83018>

[69] Yang, S.-Y., Chen, L.-Y., Najjoan, E., Kallivayalil, R.A., Viboonma, K., Jamaluddin, R., Javed, A., Hoa, D.T.Q., Iida, H., Sim, K., Swe, T., He, Y.L., Park, Y.C., Ahmed, H.U., De Alwis, A., Chiu, H.F.K., Sartorius, N., Tan, C.-H., Chong, M.Y., Shinfuku, N., Lin, S.K. (2018). Polypharmacy and Psychotropic Drug Loading in Patients with Schizophrenia in Asian Countries: The REAP-AP4 Study. *Psychiatry and Clinical Neurosciences (PCN)*.

<https://doi.org/10.1111/pcn.12676>

[68] Lin, S.-K., Lin, Y.-F., Yang, S.-Y., He, Y.-L., Kato, T., Hayakawa, K., Park, Y., Sim, K., Chiu, H.F.-K., Xiang, Y.T., Chong, M.-Y., Tan, C.-H., Inada, T., Shinfuku, N. (2018). Comparison of the Defined Daily Dose and Chlorpromazine Equivalent Methods in Antipsychotic Drug Utilization in Six Asian Countries. *Neuropsychiatry (London)*, 8, 739-744.

[67] Shinfuku, N., Hayakawa, K., Lin, S.-K., Kato, T.A. (2017). Problems of Japan's Prescription Based on the Research on East Asian Psychotropic Prescription Pattern (REAP). *Rinsyou-Seishin Yakuri (Clinical Pharmacology)*, 20, 983-991.

(In Japanese)

[full text](#)

[66] Park, Y.C., Kanba S., Chong, M.-Y., Tripathi, A., Kallivayali, Y.A., Avasth, A., Grover, S., Chee, K.Y., Tanra, A.J., Maramis, M.M., Yang, S.-Y., Sartorius, N., Tan, C-H., Shinfuku, N., Park, S.-C. (2018). Using the Brief Psychiatric Rating Scale to Detect Disorganized Speech in Schizophrenia: Findings from the REAP-AP Study. *The Kaohsiung Journal of Medical Sciences*, 34, 113-119. <https://doi.org/10.1016/j.kjms.2017.09.009>

[65] Park, S.C., Lee, M.S., Hahn, S.W., Kanba, S., Chong, M.-Y., Chee, K.Y., Udomratn, P., Tripathi, A., Sartorius, N., Shinfuku, N., Tanra, A.J., Park, Y.C. (2017). Clinical Characteristics and Psychotropic Prescribing Patterns Associated with Impaired Concentration in Asians with Depressive Disorders: The REAP-AD Study. *The Tohoku Journal of Experimental Medicine*, 242(2), 151-156.

[64] Zhong, X.M., Wang, F., Zhang, Q., Ungvari, G.S., Ng, C.H., Chiu, H.F.K., Si, T.M., Sim, K., Avasthi, A., Grover, S., Chong, M.Y., Chee, K.Y., Kanba, S., Lee, M.S., Yang, S.Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Mahendran, R., Tan, C.H., Shinfuku, N., Xiang, Y.T. (2017). Concurrent Benzodiazepine Use in Older Adults Treated with Antidepressants in Asia: International Psychogeriatrics. International Psychogeriatric Association.

[63] Rajaratnam, K., Xiang, Y.T., Tripathi, A., Chiu, H.F., Si, T.M., Chee, K.Y., Avasthi, A., Grover, S., Chong, M.Y., Kuga, H., Kanba, S., He, Y.L., Lee, M.S., Yang, S.Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Kua, E.H., Tan, C.H., Mahendran, R., Shinfuku, N., Sum, M.Y., Baldessarini, R.J., Sim, K. (2017). Clinical Use of Mood Stabilizers with Antidepressants in Asia: Report from the Research on Asian Psychotropic Prescription Patterns for Antidepressants (REAP-AD) Projects in 2004 and 2013. *Journal of Clinical Psychopharmacology*, 37, 255-259. <https://doi.org/10.1097/JCP.0000000000000670>

[62] Wang, Y.Y., Xiang, Y.T., Ungvari, G.S., Ng, C.H., Chiu, H.F.K., Yim, L.C.L., Si, T.M., Chee, K.Y., Avasthi, A., Grover, S., Chong, M.Y., Sim, K., Kanba, S., He, Y.L., Lee, M.S., Yang, S.Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M.M., Shen, W.W., Sartorius, N., Mahendran, R., Teng, J.Y., Tan, C.H., Shinfuku, N. (2017). A Comparison of Clinical Characteristics of Older Adults Treated with Antidepressants in General and Psychiatric Hospitals in Asia. *Psychogeriatrics*, 17, 348-355. <https://doi.org/10.1111/psyg.12243>

[61] Xiang, Y.T., Kato, T.A., Kishimoto, T., Ungvari, G.S., Chiu, H.F., Si, T.M., Yang, S.Y., Fujii, S., Ng, C.H., Shinfuku, N. (2017). Comparison of Treatment Patterns in Schizophrenia between China and Japan (2001-2009). *Asia Pac Psychiatry*, 9.

[60] Rajaratnam, K., Xiang, Y.T., Tripathi, A., Chiu, H.F., Si, T.M., Chee, K.Y., Avasthi, A., Grover, S., Chong, M.Y.,

Kuga, H., Kanba, S., He, Y.L., Lee, M.S., Yang, S.Y., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M., Shen, W.W., Sartorius, N., Kua, E.H., Tan, C.H., Mahendran, R., Shinfuku, N., Sum, M.Y., Baldessarini, R.J., Sim, K. (2016). Factors Associated with Antidepressant Dosing in Asia: Findings from the Research on Asian Psychotropic Prescription Study. *Journal of Clinical Psychopharmacology*, 36, 716-719. <https://doi.org/10.1097/JCP.0000000000000591>

[59] Park, S.C., Lee, M.S., Hahn, S.W., Si, T.M., Kanba, S., Chong, M.Y., Yoon, C.K., Udomratn, P., Tripathi, A., Sartorius, N., Shinfuku, N., Maramis, M.M., Park, Y.C. (2016). Suicidal Thoughts/Acts and Clinical Correlates in Patients with Depressive Disorders in Asians: Results from the REAP-AD Study. *Acta Neuropsychiatrica*, 16, 1-9.

[58] Chee, K.Y., Tripathi, A., Avasthi, A., Chong, M.Y., Xiang, Y.T., Sim, K., Kanba, S., He, Y.L., Lee, M.S., Chiu, H.F., Yang, S.Y., Kuga, H., Udomratn, P., Tanra, A.J., Maramis, M.M., Grover, S., Mahendran, R., Kallivayalil, R.A., Shen, W.W., Shinfuku, N., Tan, C.H., Sartorius, N. (2016). Prescribing Pattern of Antidepressants in Children and Adolescents: Findings from the Research on Asia Psychotropic Prescription Pattern. *East Asian Arch Psychiatry*, 26, 10-17.

[57] Wang, F., Xiang, Y.T., Ungvari, G.S., Ng, C.H., Chiu, H.F., Zheng, W., Tan, C.H., Shinfuku, N. (2016). Prevalence of Overweight in Schizophrenia Patients in Asia: Findings of the Research on Asian Psychotropic Prescription Patterns (REAP) Study. *International Journal of Clinical Pharmacology and Therapeutics*, 54, 450-455. <https://doi.org/10.5414/CP202526>

[56] Grover, S., Avasthi, A., Tripathi, A., Tanra, A.J., Chee, K.Y., He, Y.L., Chiu, H.F., Kuga, H., Lee, M.S., Chong, M.Y., Udomratn, P., Kanba, S., Yang, S.Y., Si, T.M., Sim, K., Tan, C.H., Shen, W.W., Xiang, Y.T., Sartorius, N., Shinfuku, N. (2015). Antidepressant Prescription Pattern in the Presence of Medical Co-Morbidity: REAP-AD 2013 Study. *East Asian Arch Psychiatry*, 25, 99-107.

[55] Xiang, Y.T., Ungvari, G.S., Correll, C.U., Chiu, H.F., Shinfuku, N. (2015). Trends in the Access to and the Use of Antipsychotic Medications and Psychotropic Co-Treatments in Asian Patients with Schizophrenia. *Epidemiology and Psychiatric Sciences*, 20, 1-9.

[54] Hou, C.L., Cai, M.Y., Ma, X.R., Zang, Y., Jia, F.J., Lin, Y.Q., Chiu, H.F., Ungvari, G.S., Ng, C.H., Zhong, B.L., Cao, X.L., Li, Y., Shinfuku, N., Xiang, Y.T. (2015). Clozapine Prescription and Quality of Life in Chinese Patients with Schizophrenia Treated in Primary Care. *Pharmacopsychiatry*, 48, 200-204. <https://doi.org/10.1055/s-0035-1555939>

[53] Park, S.C., Shinfuku, N., Maramis, M.M., Lee, M.S., Park, Y.C. (2015). Adjunctive Antipsychotic Prescriptions for Outpatients with Depressive Disorders in Asia: The Research on Asian Psychotropic Prescription Patterns for Antidepressants (REAP-AD) Study. *American Journal of Psychiatry*, 172, 684-685. <https://doi.org/10.1176/appi.ajp.2015.14121590>

[52] Park, S.C., Lee, M.S., Shinfuku, N., Sartorius, N., Park, Y.C. (2015). Gender Differences in Depressive Symptom Profiles and Patterns of Psychotropic Drug Usage in Asian Patients with Depression: Findings from the Research on Asian Psychotropic Prescription Patterns for Antidepressants Study. *Australian & New Zealand Journal of Psychiatry*, 0004867415579464.

<https://doi.org/10.1177/0004867415579464>

[51] Xiang, Y.T., Ungvari, G.S., Correll, C.U., Chiu, H.F., Lai, K.Y., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Yang, S.Y., Chong, M.Y., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Tan, C.H., Shinfuku, N. (2015). Use of Electroconvulsive Therapy for Asian Patients with Schizophrenia (2001-2009): Trends and Correlates. *Psychiatry and Clinical Neurosciences*, 69, 489-496.

<https://doi.org/10.1111/pcn.12283>

[50] Xiang, Y.T., Chiu, H.F., Ungvari, G.S., Correll, C.U., Lai, K.Y., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Yang, S.Y., Chong, M.Y., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Tan, C.H., Shinfuku, N. (2015). QTc Prolongation in Schizophrenia Patients in Asia: Clinical Correlates and Trends between 2004 and 2008/2009. *Human Psychopharmacology*, 30, 94-99. <https://doi.org/10.1002/hup.2458>

[49] Chen, C., Si, T.M., Xiang, Y.T., Ungvari, G.S., Wang, C.Y., He, Y.L., Kua, E.H., Fujii, S., Sim, K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Tan, C.H., Shinfuku, N. (2015). Prevalence and Prescription of Antidepressants in Depression with Somatic Comorbidity in Asia: The Research on East Asian Psychotropic Prescription Patterns Study. *Chinese Medical Journal*, 128, 853-858. <https://doi.org/10.4103/0366-6999.154272>

[48] Chee, K.Y., Tripathi, A., Avasthi, A., Chong, M.Y., Sim, K., Yang, S.Y., Glover, S., Xiang, Y.T., Si, T.M., Kanba, S., He, Y.L., Lee, M.S., Chiu, H.F., Kuga, H., Mahendran, R., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M., Shinfuku, N., Shen, W.W., Tan, C.H., Sartorius, N. (2015). International Study on Antidepressant Prescription Pattern at 40 Major Psychiatric Institutions and Hospitals in Asia: A 10-Year Comparison Study. *Asia Pacific Psychiatry*, 7, 366-374.

[47] Chee, K.Y., Tripathi, A., Avasthi, A., Chong, M.Y., Xiang, Y.T., Sim, K., Si, T.M., Kanba, S., He, Y.L., Lee, M.S., Fung-Kum Chiu, H., Yang, S.Y., Kuga, H., Udomratn, P., Kallivayalil, R.A., Tanra, A.J., Maramis, M., Grover, S., Chin, L.F., Dahlan, R., Mohamad Isa, M.F., Ebenezer, E.G., Nordin, N., Shen, W.W., Shinfuku, N., Tan, C.H., Sartorius, N. (2015). Country Variations in Depressive Symptoms Profile in Asian Countries: Findings of the Research on Asia Psychotropic Prescription (REAP) Studies. *Asia Pacific Psychiatry*, 7, 276-285. <https://doi.org/10.1111/appy.12170>

[46] Tripathi, A., Avasthi, A., Desousa, A., Bhagabati, D., Shah, N., Kallivayalil, R.A., Grover, S., Trivedi, J.K., Shinfuku, N. (2016). Prescription Pattern of Antidepressants in Five Tertiary Care Psychiatric Centres of India. *Indian Journal of Medical Research*, 143, 507-513. <https://doi.org/10.4103/0971-5916.184289>

- [45] Grover, S., Avasthi, A., Tripathi, A., Tanra, A.J., Yoon, C.K., Lin, H.Y., Chiu, H.L., Kuga, H., Soo, L.M., Chong, M.-Y., Udomratn, P., Kanba, S., Yang, S.-Y., Mei, S.T., Kang, S., Hoon, C., Shen, W., Tao, X.Y., Sartorius, N., Shinfuku, N. (2015). Antidepressant Prescription Pattern in the Presence of Medical Comorbidity: REAP-AD 2013 Study. *East Asian Archives of Psychiatry*, 25, 99-107.
- [44] Shinfuku, N. (2014). Research on Asian Prescription Patterns (REAP): Focusing on Data from Japan, Overview. *Taiwanese Journal of Psychiatry*, 28(2), 71-85.
- [43] Xiang, Y.T., Li, Y., Correll, C.U., Ungvari, G.S., Chiu, H.F., Lai, K.Y., Tang, Q.S., Hao, W., Si, T.M., Wang, C.Y., Lee, E.H., He, Y.L., Yang, S.Y., Chong, M.Y., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Tan, C.H., Shinfuku, N. (2014). Common Use of High Doses of Antipsychotic Medications in Older Asian Patients with Schizophrenia (2001-2009). *International Journal of Geriatric Psychiatry*, 29, 359-366.
<https://doi.org/10.1002/gps.4011>
- [42] Shinfuku, N. (2014). Research on Asia Psychotropic Prescription Pattern (REAP)-Overview. *Journal of Japanese Society of Social Psychiatry*, 23, 90-103. (In Japanese)
- [41] Lopez-Munoz, F., Shinfuku, N., Shen, W.W., Moreno, R., Molina, J.D., Rubio, G., Huelves, L., Noriega, C., Perez-Nieto, M.A., Alamo, C. (2013). Thirty Years of Scientific Research on Second-Generation Antipsychotic Drugs in Japan: A Bibliometric Analysis. *Open Journal of Psychiatry*, 3, 18-25. <http://www.scirp.org/journal/ojpsych/>
<https://doi.org/10.4236/ojpsych.2013.31004>
- [40] Xiang, Y.T., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., Chiu, H.F., Lai, K.Y., He, Y.L., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2013). Adjunctive Antidepressant Prescriptions for Hospitalized Patients with Schizophrenia in Asia (2001-2009). *Asia Pacific Psychiatry*, 5, E81-E87. <https://doi.org/10.1111/j.1758-5872.2012.00231.x>
- [39] Xiang, Y.T., Buchanan, R.W., Ungvari, G.S., Chiu, H.F.K., Lai, K.Y.C., Li, Y.H., Si, T.-M., Wang, C.Y., Lee, E.H.M., He, Y.L., Yang, S.-Y., Chong, M.Y., Kua, E.-H., Fujii, S., Sim, K., Yong, M.K.H., Trivedi, J.K., Chung, E.-K., Udomratn, P., Chee, K.-K., Sartorius, N., Tan, C.-H., Shinfuku, N. (2013). Use of Clozapine in Older Asian Patients with Schizophrenia between 2001 and 2009. *PLoS ONE*, 8, e66154. <https://doi.org/10.1371/journal.pone.0066154>
- [38] Xiang, Y.T., Kreyenbuhl, J., Dickerson, F.B., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., Chiu, H.F., Lai, K.Y., He, Y.L., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2012). Antipsychotic Treatment in Older Schizophrenia Patients with Extrapyrarnidal Side Effects in Asia (2001-2009). *International Journal of Clinical Pharmacology and Therapeutics*, 50, 500-504.

<https://doi.org/10.5414/CP201683>

[37] Xiang, Y.T., Dickerson, F., Kreyenbuhl, J., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., Chiu, H.F., Lai, K.Y., He, Y.L., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2012). Common Use of Anticholinergic Medications in Older Patients with Schizophrenia: Findings of the Research on Asian Psychotropic Prescription Pattern (REAP) Study, 2001-2009. *International Journal of Geriatric Psychiatry*, 28, 305-311.

[36] Xiang, Y.T., Kreyenbuhl, J., Dickerson, F.B., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2012). Use of First- and Second-Generation Antipsychotic Medications in Older Patients with Schizophrenia in Asia (2001-2009). *Australian and New Zealand Journal of Psychiatry*, 46.

[35] Xiang, Y.T., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Ungvari, G.S., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, K.H., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2012). Antipsychotic Polypharmacy in Inpatients with Schizophrenia in Asia (2001-2009). *Pharmacopsychiatry*, 45, 7-12.

<https://doi.org/10.1055/s-0031-1286345>

[34] Xiang, Y.T., Dickerson, F., Kreyenbuhl, J., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2012). Common Use of Antipsychotic Polypharmacy in Older Asian Patients with Schizophrenia (2001-2009). *Journal of Clinical Psychopharmacology*, 32, 809-813.

<https://doi.org/10.1097/JCP.0b013e3182726623>

[33] Xiang, Y.T., Dickerson, F., Kreyenbuhl, J., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2012). Adjunctive Mood Stabilizer and Benzodiazepine Use in Older Asian Patients with Schizophrenia, 2001-2009. *Pharmacopsychiatry*, 45, 217-222.

[32] Xiang, Y.T., Dickerson, F., Kreyenbuhl, J., Ungvari, G.S., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Chiu, H.F., Lai, K.Y., Shinfuku, N., Yang, S.Y., Chong, M.Y., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Tan, C.H. (2012). Prescribing Patterns of Low Doses of Antipsychotic Medications in Older Asian Patients with Schizophrenia, 2001-2009. *International Psychogeriatrics*, 24, 1002-1008.

<https://doi.org/10.1017/S1041610211002791>

[31] Ito, H., Okumura, Y., Higuchi, T., Tan, C.H., Shinfuku, N. (2012). International Variation in Antipsychotic Prescribing for Schizophrenia: Pooled Results from the Research on East Asia Psychotropic Prescription (Reap) Studies.

[30] Xiang, Y.T., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Ungvari, G.S., Chiu, H.F., Shinfuku, N., Yang, S.Y., Chong, M.Y., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Dixon, L.B., Kreyenbuhl, J.A., Tan, C.H. (2011). Clozapine Use in Schizophrenia: Findings of the Research on Asia Psychotropic Prescription (REAP) Studies from 2001 to 2009. *Australian & New Zealand Journal of Psychiatry*, 45, 968-975.

<https://doi.org/10.3109/00048674.2011.607426>

[29] Xiang, Y.T., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Ungvari, G.S., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, M.K., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2011). The Low Frequency of Reported Sexual Dysfunction in Asian Patients with Schizophrenia (2001-2009): Low Occurrence or Ignored Side Effect? *Human Psychopharmacology*, 26, 352-357.

<https://doi.org/10.1002/hup.1213>

[28] Xiang, Y.T., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Ungvari, G.S., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, K.H., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2011). Tardive Dyskinesia in the Treatment of Schizophrenia: The Findings of the Research on Asian Psychotropic Prescription Pattern (REAP) Survey (2001-2009). *International Journal of Clinical Pharmacology and Therapeutics*, 49, 382-387. <https://doi.org/10.5414/CP201505>

[27] Xiang, Y.T., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Ungvari, G.S., Chiu, H.F., Yang, S.Y., Chong, M.Y., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, K.H., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N., Shinfuku, N. (2011). Use of Anticholinergic Drugs in Patients with Schizophrenia in Asia from 2001 to 2009. *Pharmacopsychiatry*, 44, 114-118. <https://doi.org/10.1055/s-0031-1275658>

[26] Nakano, W., Fuji, S., Shinfuku, N., et al. (2011). Changes of Prescription Pattern for Schizophrenic Inpatients in Asia over the Past 10 Years: Finding from Research on Asian Psychotropic Prescription Pattern. *Rinsyou Seishinn Yakuri (Clinical Psychopharmacology)*, 14, 1397-1411. (In Japanese)

[25] Tor, P.C., Ng, T.P., Yong, K.H., Sim, K., Xiang, Y.T., Wang, C.Y., Lee, E.H., Fujii, S., Yang, S.Y., Chong, M.Y., Ungvari, G.S., Si, T., He, Y.L., Chung, E.K., Chee, K.Y., Trivedi, J., Udomratn, P., Shinfuku, N., Kua, E.H., Tan, C.H., Sartorius, N., Baldessarini, R.J. (2011). Adjunctive Benzodiazepine Treatment of Hospitalized Schizophrenia Patients in Asia from 2001 to 2008. *International Journal of Neuropsychopharmacology*, 14, 735-745.

<https://doi.org/10.1017/S146114571000163X>

[24] Sim, K., Yong, K.H., Chan, Y.H., Tor, P.C., Xiang, Y.T., Wang, C.Y., Lee, E.H., Fujii, S., Yang, S.Y., Chong, M.Y., Ungvari, G.S., Si, T., He, Y.L., Chung, E.K., Chee, K.Y., Trivedi, J., Udomratn, P., Shinfuku, N., Kua, E.H., Tan,

- C.H., Sartorius, N., Baldessarini, R.J. (2011). Adjunctive Mood Stabilizer Treatment for Hospitalized Schizophrenia Patients: Asia Psychotropic Prescription Study (2001-2008). *International Journal of Neuropsychopharmacology*, 14, 1157-1164. <https://doi.org/10.1017/S1461145711000563>
- [23] Shinfuku, N. (2011). Treatment of Schizophrenia in East Asia. In: Ishigooka, J., Okazaki, Y., and Higuchi, T., Eds., *New Strategies of Treatment of Schizophrenia*, Sentann Igakusya, Tokyo, 58-66. (In Japanese)
- [22] Xiang, Y.T., Wang, C.Y., Si, T.M., Lee, E.H., He, Y.L., Ungvari, G.S., Chiu, H.F., Yang, S.Y., Chong, M.Y., Shinfuku, N., Tan, C.H., Kua, E.H., Fujii, S., Sim, K., Yong, K.H., Trivedi, J.K., Chung, E.K., Udomratn, P., Chee, K.Y., Sartorius, N. (2010). Sex Differences in Use of Psychotropic Drugs and Drug-Induced Side Effects in Schizophrenia Patients: Findings of the Research on Asia Psychotropic Prescription (REAP) Studies. *Australian & New Zealand Journal of Psychiatry*, 45, 193-198. <https://doi.org/10.3109/00048674.2010.538839>
- [21] Nakano, W., Yang, S.-Y., Fujii, S., et al. (2010). Characteristics of Pharmacotherapy for Schizophrenic Inpatients in Japan; Findings from Research on East Asia Psychotropic Prescription Pattern of Antipsychotics. *Rinsyou Seishin Yakuri*, 13, 103-113. (In Japanese)
- [20] Sim, K., Su, H.C., Fujii, S., Yang, S.Y., Chong, M.Y., Ungvari, G., Si, T., He, Y.L., Chung, E.K., Chan, Y.H., Shinfuku, N., Kua, E.H., Tan, C.H., Sartorius, N. (2009). High-Dose Antipsychotic Use in Schizophrenia: A Comparison between the 2001 and 2004 Research on East Asia Psychotropic Prescription (REAP) Studies. *British Journal of Clinical Pharmacology*, 67, 110-117. <https://doi.org/10.1111/j.1365-2125.2008.03304.x>
- [19] Shinfuku, N. (2009). Pharmaco-Treatment of Depression in East Asia. In: Kanba, S. and Kuroki, T., Eds., *Clinics of Depression at Present*, Sougen-sya, Tokyo, 42-59. (In Japanese)
- [18] Tan, C.H., Shinfuku, N., Sim, K. (2008). Psychotropic Prescription Practices in East Asia: Looking Back and Peering Ahead. *Current Opinion in Psychiatry*, 21, 645-650. <https://doi.org/10.1097/YCO.0b013e32830e6dc4>
- [17] Shinfuku, N., Tan, C.H. (2008). Pharmacotherapy for Schizophrenic Inpatients in East Asia—Changes and Challenges. *International Review of Psychiatry*, 20, 460-468. <https://doi.org/10.1080/09540260802397560>
- [16] Sim, K., Chuan, S.H., Fujii, S., Yang, S.Y., Chong, M.Y., Si, T., He, Y.L., Chung, E.K., Chan, Y.H., Shinfuku, N., Hoon Tan, C., Ungvari, G., Baldessarini, R.J. (2008). Low Doses of Antipsychotic Drugs for Hospitalized Schizophrenia Patients in East Asia: 2004 vs. 2001. *International Journal of Neuropsychopharmacology*, 12, 117-123. <https://doi.org/10.1017/S1461145708009280>
- [15] Uchida, N., Chong, M.Y., Tan, C.H., Nagai, H., Tanaka, M., Lee, M.S., Fujii, S., Yang, S.Y., Si, T., Sim, K., Wei,

H., Ling, H.Y., Nishimura, R., Kawaguchi, Y., Edwards, G., Sartorius, N., Shinfuku, N. (2007). International Study on Antidepressant Prescription Pattern at 20 Teaching Hospitals and Major Psychiatric Institutions in East Asia: Analysis of 1898 Cases from China, Japan, Korea, Singapore and Taiwan. *Psychiatry and Clinical Neurosciences*, 61, 522-528.

<https://doi.org/10.1111/j.1440-1819.2007.01702.x>

[14] Tanaka, M., Nagai, H., Uchida, N., et al. (2007). Current Situation on the Prescription of Antidepressants in East Asia-Findings from REAP Collaborative Surveys in Five Countries and Areas. *Rinsyouseishin Yakuri (Clinical Psychopharmacology)*, 10, 131-146. (In Japanese)

[13] Sim, K., Lee, N.B., Chua, H.C., Mahendran, R., Fujii, S., Yang, S.Y., Chong, M.Y., Si, T., He, Y.L., Lee, M.S., Sung, K.M., Chung, E.K., Chan, Y.H., Shinfuku, N., Tan, C.H., Sartorius, N., Baldessarini, R.J. (2007). Newer Antidepressant Drug Use in East Asian Psychiatric Treatment Settings: REAP (Research on East Asia Psychotropic Prescriptions) Study. *British Journal of Clinical Pharmacology*, 63, 431-437. <https://doi.org/10.1111/j.1365-2125.2006.02780.x>

[12] Yoshimura, R., Okamoto, T., Nakamura, J., Tateno, M., Otsuka, K., Takahashi, H., Fujisawa, D., Takamatsu, T., Fujii, S., Sato, S., Inoue, M., Sasaki, H., Kuroki, T., Shinfuku, N. (2006). Prescription Pattern of Antipsychotic Drugs for Schizophrenic Inpatients in Japan: Research on East Asia Psychotropic Prescription Pattern-Antipsychotics study. *Psychiatry and Clinical Neurosciences*, 60, 778-779. <https://doi.org/10.1111/j.1440-1819.2006.01601.x>

[11] Fujii, S., Kaneko, N., Kitsukawa, H., et al. (2005). Prescription of Antidepressants in Japan—Finding from International Collaborative Study on the Prescription of Antidepressants in Five Countries and Areas in Asia. *Japanese Journal of Social Psychiatry*, 14, 30-35. (In Japanese)

[10] Fujii, S., Shinfuku, N. (2005). Historical Overview on the Use of Levomepromazine in Japan and the Setting of Its Dose in Pharmacotherapy. *Rinsyouseishin Yakuri (Japanese Journal of Clinical Psychiatry)*, 8, 1227-1238. (In Japanese)

[9] Fujii, S., Shinfuku, N. (2004). The Characteristics of Pharmacotherapy for the Inpatients with Schizophrenia in Japan—Comparison with Other Countries and Regions. *Rinsyouseishin Yakuri (Japanese Journal of Clinical Psychiatry)*, 7,3-14. (In Japanese)

[8] Sim, K., Su, A., Chan, Y.H., Shinfuku, N., Kua, E.H., Tan, C.H. (2004). Clinical Correlates of Antipsychotic Polytherapy in Patients with Schizophrenia in Singapore. *Psychiatry and Clinical Neurosciences*, 58, 324-329. <https://doi.org/10.1111/j.1440-1819.2004.01239.x>

[7] Sim, K., Su, A., Ungvari, G.S., Fujii, S., Yang, S.Y., Chong, M.Y., Si, T., Chung, E.K., Tsang, H.Y., Chan, Y.H., Shinfuku, N., Tan, C.H. (2004). Depot Antipsychotic Use in Schizophrenia: An East Asian Perspective. *Human*

Psychopharmacology, 19, 103-109. <https://doi.org/10.1002/hup.571>

[6] Sim, K., Su, A., Leong, J.Y., Yip, K., Chong, M.Y., Fujii, S., Yang, S., Ungvari, G.S., Si, T., Chung, E.K., Tsang, H.Y., Shinfuku, N., Kua, E.H., Tan, C.H. (2004). High Dose Antipsychotic Use in Schizophrenia: Findings of the REAP (Research on East Asia Psychotropic Prescriptions) Study. *Pharmacopsychiatry*, 37, 175-179. <https://doi.org/10.1055/s-2004-827174>

[5] Sim, K., Su, A., Fujii, S., Yang, S.Y., Chong, M.Y., Ungvari, G.S., Si, T., Chung, E.K., Tsang, H.Y., Chan, Y.H., Heckers, S., Shinfuku, N., Tan, C.H. (2004). Antipsychotic Polypharmacy in Patients with Schizophrenia: A Multicentre Comparative Study in East Asia. *British Journal of Clinical Pharmacology*, 58, 178-183. <https://doi.org/10.1111/j.1365-2125.2004.02102.x>

[4] Chong, M.Y., Tan, C.H., Fujii, S., Yang, S.Y., Ungvari, G.S., Si, T., Chung, E.K., Sim, K., Tsang, H.Y., Shinfuku, N. (2004). Antipsychotic Drug Prescription for Schizophrenia in East Asia: Rationale for Change. *Psychiatry and Clinical Neurosciences*, 58, 61-67. <https://doi.org/10.1111/j.1440-1819.2004.01194.x>

[3] Si, T.M., He, Y.L., Zhou, R.Y., et al. (2003). Psychotropic Prescription Pattern for the Inpatients with Schizophrenia. *Journal of Chinese Mental Health*, 17, 705-707. (In Chinese)

[2] Fujii, S., Maeda, K., Shinfuku, N. (2003). International Collaborative Studies on the Prescription of Antipsychotics in East Asia—Finding from Research on East Asia Psychotropic Prescription Pattern. *Rinsyou Seishin Igaku (Clinical Psychiatry)*, 32, 629-646. (In Japanese)

[1] Tan, C.H. (2000) New Challenges in Psychopharmacology. *Singapore Med. J*, 41, 3-4.